
Limahuli, Kaua'i



WATERSHED FEATURES

Limahuli watershed occurs on the island of Kaua'i. The Hawaiian meaning of the name is "turned hand". The area of the watershed is 1.8 square mi (4.7 square km), with maximum elevation of 3287 ft (1002 m). The watershed's DAR cluster code is 2, meaning that the watershed is small, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 0% agricultural, 100% conservation, 0% rural, and 0% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	County	Nature Conservancy	Other Private
0.0	0.0	75.7	8.6	0.0	0.0	15.7

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity	Managed for Multiple	Protected but	
<u>Protection</u>	<u>Uses</u>	<u>Unmanaged</u>	<u>Unprotected</u>
0.0	0.0	75.7	24.3

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	Square mi	Square km
High Intensity Developed	0.0	0.00	0.00
Low Intensity Developed	0.9	0.02	0.04
Cultivated	0.0	0.00	0.00
Grassland	0.9	0.02	0.04
Scrub/Shrub	78.4	1.42	3.67
Evergreen Forest	18.3	0.33	0.86
Palustrine Forested	0.1	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	1.3	0.02	0.06
Unconsolidated Shoreline	0.0	0.00	0.00
Water	0.1	0.00	0.01
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Limahuli is a perennial stream. Total stream length is 4.8 mi (7.8 km). The terminal stream order is 2.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	Lower	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	0.0	26.2	52.1	21.7

The following stream(s) occur in the watershed: Limahuli

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1966 1979 1981 1989 1990 2001 2003

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

Survey type	<u>Estuary</u>	Lower	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	0	2	0	0
HDFG	0	0	2	0	0
Published Report	0	1	4	1	0
USGS Surveys	0	0	1	0	0

BIOTA INFORMATION

Species List

Native Specie	s	Native Species		
Crustaceans	Amphipod sp.	Insects	Empidid sp.	
	Atyoida bisulcata		Limonia sp.	
	Macrobrachium grandimanus		Megalagrion vagabundum	
Fish	Awaous guamensis		Orthocladius sp.	
	Eleotris sandwicensis		Saldula exulans	
	Gobiid sp.		Telmatogeton sp.	
	Kuhlia xenura		Tipulid sp.	
	Lentipes concolor			
	Sicyopterus stimpsoni			
Snails	Neritina granosa			
Worms	Hirudinean sp.			
	Oligochaete sp.			

Introduced Species Introduced Species

Crustaceans	lsopod sp. <i>Macrobrachium lar</i>	Insects	Cheumatopsyche analis Chironomid larvae
Snails	Lymnaeid sp.		Hydroptila potosina
Worms	Dugesia sp.		Ischnura posita
			Trichoptera larvae

Species Distributions: Presence (P) of species in different stream reaches.

Scientific Name	<u>Status</u>	<u>Estuary</u>	Lower	<u>Middle</u>	Upper Headwaters
Atyoida bisulcata	Endemic			Р	Р
Macrobrachium grandimanus	Endemic		Р		
Eleotris sandwicensis	Endemic		Р		
Kuhlia xenura	Endemic		Р		
Lentipes concolor	Endemic			Р	
Sicyopterus stimpsoni	Endemic		Р	Р	
Megalagrion vagabundum	Endemic			Р	
Orthocladius sp.	Endemic			Р	
Saldula exulans	Endemic				Р

Neritina granosa	Endemic		Р		Р
Amphipod sp.	Indigenous		Р		
Awaous guamensis	Indigenous		Р		
Gobiid sp.	Indigenous		Р		
Limonia sp.	Indigenous		Р		
Telmatogeton sp.	Indigenous		Р		
Isopod sp.	Introduced		Р		
Macrobrachium lar	Introduced	Р	Р		
Cheumatopsyche analis	Introduced		Р		Р
Chironomid larvae	Introduced		Р		Р
Hydroptila potosina	Introduced		Р		Р
Ischnura posita	Introduced		Р		
Trichoptera larvae	Introduced		Р		
Lymnaeid sp.	Introduced		Р		
Dugesia sp.	Introduced		P F)	
Empidid sp.	Undetermined		Р		
Tipulid sp.	Undetermined		Р		Р
Hirudinean sp.	Undetermined		Р		
Oligochaete sp.	Undetermined		Р		

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Substantial

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity > 19 spp.	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 Introduced
No	Yes	Yes
Abundance of Any Native Species	Presence of Candidate Endangered Species	Endangered Newcomb's Snail Habitat
Yes	No	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Limahuli, Kaua'i


<u>Land Cover Rating</u>: Rating is based on a scoring sytem where in general forested lands score positively and developed lands score negatively.



<u>Shallow Waters Rating</u>: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.



<u>Stewardship Rating</u>: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.





WATERSHED RATING (Cont): Limahuli, Kaua'i

<u>Size Rating</u>: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.



2







Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.

Watershed Rating

4







<u>Reach Diversity Rating</u>: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.

Watershed Rating

5







<u>Total Watershed Rating</u>: Rating is based on combination of <u>Land Cover Rating</u>, <u>Shallow Waters Rating</u>, <u>Stewardship Rating</u>, <u>Size Rating</u>, <u>Wetness Rating</u>, and <u>Reach Diversity Rating</u>.

Watershed Rating

5







BIOLOGICAL RATING: Limahuli, Kaua'i

<u>Native Species Rating</u>: Rating is based on the number of native species observed in the watershed.

Stream Rating

8







<u>Introduced Genera Rating</u>: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating

10







<u>All Species' Score Rating:</u> Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating

6







<u>Total Biological Rating</u>: Rating is the combination of the <u>Native Species Rating</u>, <u>Introduced Genera Rating</u>, and the <u>All Species' Score Rating</u>.

Stream Rating


8






OVERALL RATING: Limahuli, Kaua'i

Overall Rating: Rating is a combination of the <u>Total Watershed Rating</u> and the <u>Total Biological Rating</u>.



RATING STRENGTH: Limahuli, Kaua'i

<u>Rating Strength:</u> Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

- 1967. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
- 1991. Honigman, L. and A. Newman. A Biological Database of Aquatic Resources on Hawaiian Streams. Proceedings of the 1990 Symposium on Freshwater Stream Biology and Fisheries Management. 51-76.
- 2005. Englund, R.A., Laederich, H.M., Ghotaslou, A., Bollick, H. and A.A. Scott. Analysis of Kaua'i Aquatic Invertebrate Benthic Samples. Final Report, Hawaii Biological Survey.
- 2005. USGS. Stream Quality Indicators of Hawai'i.

- 2006. Brasher, A.M.D., Lunton, C.D., Goodbred, S.L. and R.H. Wolff. Invasion Patterns Along Elevation and Urbanization Gradients in Hawaiian Streams. Transactions of the American Fisheries Society. 135. 1109-1129.
- 2006. Polhemus, D.A. Maps of Damselfly Locations.
- 2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
- 2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.

Blank Page