


Wailua River, Kaua'i



WATERSHED FEATURES

Wailua River watershed occurs on the island of Kaua'i. The Hawaiian meaning of the name is “two waters”. The area of the watershed is 52.6 square mi (136.2 square km), with maximum elevation of 5226 ft (1593 m). The watershed’s DAR cluster code is 8, meaning that the watershed is very large, and steep in the upper watershed.. The percent of the watershed in the different land use districts is as follows: 27.8% agricultural, 69% conservation, 1.4% rural, and 1.8% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other</u>	<u>Private</u>
0.0	0.0	69.3	0.0	0.1	0.0		30.6

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
0.0	0.0	69.0	31.0

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	0.0	0.02	0.04
Low Intensity Developed	1.7	0.90	2.32
Cultivated	6.1	3.19	8.27
Grassland	8.5	4.48	11.61
Scrub/Shrub	39.1	20.58	53.29
Evergreen Forest	40.7	21.38	55.37
Palustrine Forested	0.3	0.15	0.39
Palustrine Scrub/Shrub	0.6	0.30	0.78
Palustrine Emergent	0.1	0.07	0.17
Estuarine Forested	0.0	0.00	0.00
Bare Land	0.0	0.03	0.06
Unconsolidated Shoreline	0.0	0.00	0.01
Water	0.4	0.23	0.60
Unclassified	2.4	1.27	3.28

STREAM FEATURES

Wailua River is a perennial stream. Total stream length is 177.3 mi (285.4 km). The terminal stream order is 5.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
1.4	0.0	23.6	72.8	2.2

The following stream(s) occur in the watershed:

Hāli'i	'Ili'i'i'ula	'Iole	Kalama	Kaulu
Kāwī	North Fork	'Ōpaeka'a	Palikea	South Fork
Uhau'i'ole	Waiahi	Wai'aka	Waikoko	Wailua

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1962	1963	1977	1978	1986	1990	1994
2003	2004					

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	2	0	26	0
DAR Observation	0	0	2	3	0
DAR Point Quadrat	0	0	53	27	0
HDFG	3	3	6	23	0
Published Report	0	1	2	3	0
Reservoir	0	0	4	1	0

BIOTA INFORMATION**Species List****Native Species**

Crustaceans	Amphipod sp. <i>Atyoida bisulcata</i> Copepod sp. <i>Macrobrachium grandimanus</i> Ostracod sp.
Fish	<i>Awaous guamensis</i> <i>Chanos chanos</i> <i>Eleotris sandwicensis</i> <i>Elops hawaiiensis</i> Gobiid sp. <i>Kuhlia sandwicensis</i> <i>Kuhlia</i> sp. <i>Mugil cephalus</i> Mugilid sp. <i>Sicyopterus stimpsoni</i>
Worms	<i>Namalycastis</i> sp.

Native Species

Insects	<i>Anax junius</i> <i>Anax</i> sp. <i>Campsicnemus bicoloripes</i> Ephydrid sp. <i>Megalagrion eudytum</i> <i>Megalagrion heterogamias</i> <i>Megalagrion oresitrophum</i> <i>Megalagrion orobates</i> <i>Megalagrion</i> sp. <i>Megalagrion vagabundum</i> <i>Scatella cilipes</i> <i>Telmatogeton</i> sp. Tipulid sp.
----------------	---

Introduced Species

Clams	<i>Corbicula fluminea</i>
Crustaceans	<i>Macrobrachium lar</i> <i>Procambarus clarkii</i>
Fish	<i>Clarias fuscus</i> <i>Cyprinus carpio</i> <i>Gambusia affinis</i> <i>Lepomis macrochirus</i> <i>Lepomis</i> sp. <i>Micropterus dolomieu</i> <i>Micropterus salmoides</i> <i>Micropterus</i> sp. <i>Oreochromis mossambicus</i>

Introduced Species

Insects	<i>Cheumatopsyche analis</i> Chironomid larvae Culicid sp. <i>Enallagma civile</i> <i>Ischnura posita</i> <i>Ischnura ramburi</i>
----------------	--

	<i>Poecilia reticulata</i>
	<i>Tilapia sp.</i>
	unidentified cyprinid
	<i>Xiphophorus helleri</i>
Snails	<i>Helisoma sp.</i>
	Lymnaeid sp.
	<i>Tarebia granifera</i>
	Thiarid sp.
Worms	<i>Dugesia sp.</i>

Species found in Impoundments

Fish	<i>Clarias fuscus</i>
	<i>Lepomis sp.</i>
	<i>Micropterus sp.</i>
	<i>Tilapia sp.</i>
	unidentified cyprinid

Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
<i>Corbicula fluminea</i>	Introduced	0.5	1	0.6
<i>Atyoida bisulcata</i>	Endemic	0.25	0.25	0.3
<i>Macrobrachium lar</i>	Introduced	2	2	2.0
<i>Awaous guamensis</i>	Indigenous	1	2.75	1.6
Gobiid sp.	Indigenous	0.75	0.75	0.8
<i>Micropterus dolomieu</i>	Introduced	8	15	10.8
<i>Tarebia granifera</i>	Introduced	0.25	1	0.5

Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic				0.06	
<i>Awaous guamensis</i>	Indigenous			0.27		
<i>Corbicula fluminea</i>	Introduced			0.17	0.18	
<i>Macrobrachium lar</i>	Introduced			0.02		
<i>Micropterus dolomieu</i>	Introduced			0.07	0.06	
<i>Tarebia granifera</i>	Introduced				7.69	

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic		P	P	P	
<i>Macrobrachium grandimanus</i>	Endemic	P	P			
<i>Eleotris sandwicensis</i>	Endemic	P	P			
<i>Sicyopterus stimpsoni</i>	Endemic		P			
<i>Campsicnemus bicoloripes</i>	Endemic			P		

Wailua River, Kaua'i

<i>Megalagrion eudytum</i>	Endemic				P		
<i>Megalagrion heterogamias</i>	Endemic				P		
<i>Megalagrion oresitrophum</i>	Endemic				P		
<i>Megalagrion orobates</i>	Endemic				P		
<i>Megalagrion sp.</i>	Endemic	P	P	P	P		
<i>Megalagrion vagabundum</i>	Endemic		P		P		
<i>Scatella cilipes</i>	Endemic			P			
Amphipod sp.	Indigenous					P	
<i>Awaous guamensis</i>	Indigenous	P	P	P	P		
<i>Chanos chanos</i>	Indigenous	P	P				
<i>Elops hawaiiensis</i>	Indigenous	P					
Gobiid sp.	Indigenous	P	P	P	P		
<i>Kuhlia sandvicensis</i>	Indigenous	P	P				
<i>Kuhlia sp.</i>	Indigenous	P					
<i>Mugil cephalus</i>	Indigenous	P	P				
Mugilid sp.	Indigenous	P					
<i>Anax junius</i>	Indigenous					P	
<i>Anax sp.</i>	Indigenous					P	
<i>Telmatogeton sp.</i>	Indigenous		P	P	P		
<i>Namalycastis sp.</i>	Indigenous			P			
<i>Corbicula fluminea</i>	Introduced			P	P		
<i>Macrobrachium lar</i>	Introduced			P	P	P	P
<i>Procambarus clarkii</i>	Introduced	P	P	P	P		
<i>Clarias fuscus</i>	Introduced		P	P	P		
<i>Cyprinus carpio</i>	Introduced		P		P		
<i>Gambusia affinis</i>	Introduced				P		
<i>Lepomis macrochirus</i>	Introduced	P	P	P	P		
<i>Lepomis sp.</i>	Introduced	P		P	P		
<i>Micropterus dolomieu</i>	Introduced	P	P	P	P		
<i>Micropterus salmoides</i>	Introduced		P	P	P		
<i>Micropterus sp.</i>	Introduced			P	P		
<i>Oreochromis mossambicus</i>	Introduced	P					
<i>Poecilia reticulata</i>	Introduced		P		P		
<i>Tilapia sp.</i>	Introduced	P		P	P		
unidentified cyprinid	Introduced			P	P		

<i>Xiphophorus helleri</i>	Introduced	P	P	P
<i>Cheumatopsyche analis</i>	Introduced		P	P
Chironomid larvae	Introduced		P	P
Culicid sp.	Introduced			P
<i>Enallagma civile</i>	Introduced			P
<i>Ischnura posita</i>	Introduced	P		P
<i>Ischnura ramburi</i>	Introduced			P
<i>Helisoma</i> sp.	Introduced			P
Lymnaeid sp.	Introduced	P		P
<i>Tarebia granifera</i>	Introduced			P
Thiarid sp.	Introduced			P
<i>Dugesia</i> sp.	Introduced			P
Copepod sp.	Undetermined		P	
Ostracod sp.	Undetermined		P	
Ephydrid sp.	Undetermined			P
Tipulid sp.	Undetermined			P

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Moderate

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity <u>> 19 spp.</u>	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 <u>Introduced</u>
No	Yes	No
Abundance of Any <u>Native Species</u>	Presence of Candidate <u>Endangered Species</u>	Endangered Newcomb's <u>Snail Habitat</u>
No	No	Yes

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Wailua River, Kaua'i


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




WATERSHED RATING (Cont): Wailua River, Kaua'i


Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.





Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.



BIOLOGICAL RATING: Wailua River, Kaua'i



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream Rating
7





Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating
3





All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating
3



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.


Stream Rating
3



OVERALL RATING: Wailua River, Kaua'i

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.


Watershed
Rating
7



RATING STRENGTH: Wailua River, Kaua'i

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.

Information
Rating
7



REFERENCES

1963. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.

1964. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.

1986. Kahala Energy Development Corporation. Wailuku River Hydropower Kaua'i, Hawai'i. Final Environmental Impact Statement.

1990. Englund, R.A., Ryel, R., and V. Lamarra. Habitat selection of 'O'opu Nakea and exotic stream fishes in the South Fork Wailua River and Hanalei River, Kaua'i County, Hawai'i.

1992. Baker, J.A. and S.A. Foster. Estimating Density and Abundance of Endemic Fishes in Hawaiian Streams.
1994. Way, C.M. Proceedings of the International Symposium on Hawaiian Stream Ecology, Preservations, and Management. U.S. Army Corps of Engineers.
2005. Englund, R.A., Laederich, H.M., Ghotaslou, A., Bollick, H. and A.A. Scott. Analysis of Kaua'i Aquatic Invertebrate Benthic Samples. Final Report, Hawaii Biological Survey.
2006. Polhemus, D.A. Maps of Damselfly Locations.
2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.
2008. Hawai'i Division of Aquatic Resources. Impoundment Surveys in DAR Aquatic Surveys Database.

Blank Page