


# Honokōhau, Maui



## WATERSHED FEATURES

Honokōhau watershed occurs on the island of Maui. The Hawaiian meaning of the name is “bay drawing dew”. The area of the watershed is 8.1 square mi (21 square km), with maximum elevation of 5778 ft (1761 m). The watershed's DAR cluster code is 6, meaning that the watershed is large, narrow, and steep in the upper watershed. The percent of the watershed in the different land use districts is as follows: 6% agricultural, 94% conservation, 0% rural, and 0% urban.

**Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.**

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other</u>	<u>Private</u>
0.0	0.0	2.4	0.0	0.0	0.0		97.6

**Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.**

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
74.8	2.7	0.0	22.5

**Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.**

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	0.0	0.00	0.00
Low Intensity Developed	0.1	0.00	0.01
Cultivated	1.3	0.10	0.27
Grassland	0.8	0.06	0.17
Scrub/Shrub	13.9	1.13	2.92
Evergreen Forest	83.3	6.75	17.48
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.3	0.03	0.07
Estuarine Forested	0.0	0.00	0.00
Bare Land	0.2	0.02	0.04
Unconsolidated Shoreline	0.1	0.01	0.02
Water	0.0	0.00	0.01
Unclassified	0.0	0.00	0.00

### STREAM FEATURES

Honokōhau is a perennial stream. Total stream length is 20.2 mi (32.4 km). The terminal stream order is 2.

**Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.**

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	4.0	21.0	52.3	22.7

The following stream(s) occur in the watershed:  
Honokōhau

### BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1961      1990      1992      1997      2003      2004

**Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.**

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	0	0	1	5
DAR General Surveys	0	1	0	0	0
DAR Observation	0	2	2	0	0
DAR Point Quadrat	0	49	15	50	0
HDFG	0	0	2	2	0
Published Report	0	0	0	2	0

**BIOTA INFORMATION**

**Species List**

**Native Species**

<b>Crustaceans</b>	<i>Atyoida bisulcata</i>
<b>Fish</b>	<i>Awaous guamensis</i>
	<i>Eleotris sandwicensis</i>
	Gobiid sp.
	<i>Kuhlia</i> sp.
	<i>Kuhlia xenura</i>
	<i>Lentipes concolor</i>
	<i>Sicyopterus stimpsoni</i>
<b>Snails</b>	<i>Neritina granosa</i>

**Native Species**

<b>Insects</b>	<i>Anax junius</i>
	<i>Anax strenuus</i>
	<i>Campsicnemus</i> sp.
	<i>Chironomus hawaiiensis</i>
	<i>Dasyhelea hawaiiensis</i>
	<i>Eurynogaster</i> sp.
	<i>Hyposmocoma</i> sp.
	<i>Limonia jacobus</i>
	<i>Megalagrion blackburni</i>
	<i>Megalagrion calliphya</i>
	<i>Megalagrion hawaiiense</i>
	<i>Megalagrion nigrohamatum</i>
	<i>nigrohamatum</i>
	<i>Megalagrion pacificum</i>
	<i>Megalagrion</i> sp.
	<i>Paraliancalus metallicus</i>
	<i>Procanacae acuminata</i>
	<i>Procanace confusa</i>
	<i>Procanace constricta</i>
	<i>Saldula exulans</i>
	<i>Scatella cilipes</i>
	<i>Scatella clavipes</i>
	<i>Scatella femoralis</i>
	<i>Scatella mauiensis</i>
	<i>Scatella oahuense</i>
	<i>Scatella warreni</i>
	<i>Sigmataneurum</i> sp.
	<i>Telmatogeton</i> sp.
	<i>Telmatogeton torrenticola</i>

**Introduced Species**

<b>Crustaceans</b>	<i>Macrobrachium lar</i> <i>Procambarus clarkii</i>
<b>Fish</b>	<i>Hypostomus watwata</i> <i>Poecilia reticulata</i> Poeciliid sp.
<b>Snails</b>	<i>Melania sp.</i>

**Introduced Species**

<b>Insects</b>	<i>Cheumatopsyche analis</i> Chironomid larvae <i>Cricotopus bicinctus</i> <i>Hydroptila potosina</i>
----------------	--

**Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.**

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
<i>Atyoida bisulcata</i>	Endemic	1.25	2	1.9
<i>Macrobrachium lar</i>	Introduced	1.25	1.25	1.3
<i>Eleotris sandwicensis</i>	Endemic	3	3.25	3.1
<i>Kuhlia xenura</i>	Endemic	0.75	3	1.6
<i>Lentipes concolor</i>	Endemic	2	5	3.5
<i>Sicyopterus stimpsoni</i>	Endemic	2.5	5	3.5
<i>Awaous guamensis</i>	Indigenous	3	8	4.2
<i>Kuhlia sp.</i>	Indigenous	1	1.25	1.2
Poeciliid sp.	Introduced	0.75	1	0.9
<i>Anax junius</i>	Indigenous	0.75	0.75	0.8

**Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.**

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic				11.7	
<i>Eleotris sandwicensis</i>	Endemic		0.07			
<i>Kuhlia xenura</i>	Endemic		0.81			
<i>Lentipes concolor</i>	Endemic				0.44	
<i>Megalagrion blackburni</i>	Endemic				0.01	
<i>Megalagrion nigrohamatum</i>	Endemic				0.01	
<i>Sicyopterus stimpsoni</i>	Endemic		0.07	0.33		
<i>Anax junius</i>	Indigenous				0.01	
<i>Awaous guamensis</i>	Indigenous		0.37	0.44		
<i>Kuhlia sp.</i>	Indigenous		0.1			
<i>Telmatogeton sp.</i>	Indigenous				0.01	
<i>Macrobrachium lar</i>	Introduced		0.03			
Poeciliid sp.	Introduced		0.1			

**Species Distributions: Presence (P) of species in different stream reaches.**

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic		P	P	P	
<i>Eleotris sandwicensis</i>	Endemic		P			
<i>Kuhlia xenura</i>	Endemic		P			
<i>Lentipes concolor</i>	Endemic		P		P	

<i>Sicyopterus stimpsoni</i>	Endemic	P	P	P	
<i>Anax strenuus</i>	Endemic			P	
<i>Chironomus hawaiiensis</i>	Endemic			P	
<i>Dasyhelea hawaiiensis</i>	Endemic			P	
<i>Hyposmocoma sp.</i>	Endemic			P	
<i>Limonia jacobus</i>	Endemic			P	
<i>Megalagrion blackburni</i>	Endemic			P	
<i>Megalagrion calliphya</i>	Endemic			P	P
<i>Megalagrion hawaiiense</i>	Endemic			P	P
<i>Megalagrion nigrohamatum nigrohamatum</i>	Endemic			P	P
<i>Megalagrion pacificum</i>	Endemic			P	
<i>Megalagrion sp.</i>	Endemic		P	P	
<i>Paraliancalus metallicus</i>	Endemic			P	
<i>Procanacae acuminata</i>	Endemic			P	
<i>Procanace confusa</i>	Endemic			P	
<i>Procanace constricta</i>	Endemic			P	
<i>Saldula exulans</i>	Endemic			P	
<i>Scatella cilipes</i>	Endemic			P	
<i>Scatella clavipes</i>	Endemic			P	
<i>Scatella femoralis</i>	Endemic			P	
<i>Scatella mauiensis</i>	Endemic			P	
<i>Scatella oahuense</i>	Endemic			P	
<i>Scatella warreni</i>	Endemic			P	
<i>Telmatogeton torrenticola</i>	Endemic			P	
<i>Neritina granosa</i>	Endemic			P	
<i>Awaous guamensis</i>	Indigenous	P	P		
Gobiid sp.	Indigenous		P	P	
<i>Kuhlia sp.</i>	Indigenous	P			
<i>Anax junius</i>	Indigenous			P	
<i>Campsicnemus sp.</i>	Indigenous			P	
<i>Eurynogaster sp.</i>	Indigenous			P	
<i>Sigmataneurum sp.</i>	Indigenous			P	
<i>Telmatogeton sp.</i>	Indigenous		P	P	
<i>Macrobrachium lar</i>	Introduced	P			
<i>Procambarus clarkii</i>	Introduced	P			
<i>Poecilia reticulata</i>	Introduced	P			
Poeciliid sp.	Introduced	P			
<i>Cheumatopsyche analis</i>	Introduced			P	
Chironomid larvae	Introduced	P	P	P	
<i>Cricotopus bicinctus</i>	Introduced			P	
<i>Hydroptila potosina</i>	Introduced			P	
<i>Melania sp.</i>	Introduced		P	P	

## HISTORIC RANKINGS

**Historic Rankings:** These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Outstanding

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

**Current DAR Decision Rule Status:** The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.

Native Insect Diversity  
> 19 spp.

Yes

Native Macrofauna  
Diversity > 5 spp.

Yes

Absence of Priority 1  
Introduced

No

Abundance of Any  
Native Species

No

Presence of Candidate  
Endangered Species

Yes

Endangered Newcomb's  
Snail Habitat


No

### CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

#### WATERSHED RATING: Honokōhau, Maui


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




### WATERSHED RATING (Cont): Honokōhau, Maui


Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.





Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.



### BIOLOGICAL RATING: Honokōhau, Maui



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream Rating  
**8**





Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating  
**9**





All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating  
**9**



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.


Stream Rating  
**8**



### OVERALL RATING: Honokōhau, Maui

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.


Watershed  
Rating  
**9**



### RATING STRENGTH: Honokōhau, Maui

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.

Information  
Rating  
**7**



### REFERENCES

- 1961. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
- 1990. Hau, S. Memo Notebook: 1990 FW Stream Surveys Vol. 2.
- 1997. Tagawa, A.W. Management of a Database on the Occurrence, Abundance and Distribution of Native Freshwater Species. Job Progress Report.
- 2003. Englund, R.A. et al. Systematic Inventory of Rare and Alien Aquatic Species in Selected O‘ahu, Maui, and Hawai‘i island Streams. Hawaii Biological Survey.
- 2006. Polhemus, D.A. Maps of Damselfly Locations.

2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.

2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.

Blank Page