


# Waiehu, Maui



## WATERSHED FEATURES

Waiehu watershed occurs on the island of Maui. The Hawaiian meaning of the name is “water spray”. The area of the watershed is 5.2 square mi (13.4 square km), with maximum elevation of 5161 ft (1573 m). The watershed’s DAR cluster code is not yet determined. The percent of the watershed in the different land use districts is as follows: 24.8% agricultural, 59.4% conservation, 1.5% rural, and 14.3% urban.

**Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.**

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other</u>	<u>Private</u>
1.6	0.0	1.5	0.1	0.0	0.0		96.9

**Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.**

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
0.0	46.2	2.8	50.9

**Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.**

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	2.6	0.13	0.34
Low Intensity Developed	7.7	0.40	1.03
Cultivated	20.8	1.08	2.79
Grassland	2.3	0.12	0.32
Scrub/Shrub	14.8	0.77	1.99
Evergreen Forest	50.0	2.60	6.72
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	1.2	0.06	0.16
Unconsolidated Shoreline	0.2	0.01	0.03
Water	0.4	0.02	0.05
Unclassified	0.0	0.00	0.00

**STREAM FEATURES**

Waiehu is a perennial stream. Total stream length is 14.7 mi (23.7 km). The terminal stream order is 3.

**Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.**

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	4.2	39.8	50.0	6.1

The following stream(s) occur in the watershed:  
 North Waiehu    South Waiehu    Waiehu

**BIOTIC SAMPLING EFFORT**

Biotic samples were gathered in the following year(s):

1961	1962	1976	1990	1991	1994	1995
2000	2005					

**Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.**

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	0	1	0	0
DAR General Surveys	0	1	0	0	0
DAR Observation	0	3	0	0	0
DAR Point Quadrat	0	27	0	0	0
HDFG	0	0	3	3	0
Published Report	0	1	0	0	0
Reservoir	0	0	3	0	0

**BIOTA INFORMATION**

**Species List**

**Native Species**

<b>Crustaceans</b>	<i>Atyoida bisulcata</i>
<b>Fish</b>	<i>Awaous guamensis</i>
	<i>Eleotris sandwicensis</i>
	Gobiid sp.
	<i>Kuhlia sandwicensis</i>
	<i>Kuhlia sp.</i>
	<i>Kuhlia xenura</i>
	<i>Sicyopterus stimpsoni</i>
<b>Snails</b>	<i>Clithon cariosus</i>
	<i>Neritina granosa</i>
	<i>Neritina vespertina</i>

**Native Species**

<b>Insects</b>	<i>Anax junius</i>
	<i>Campsicnemus ridiculus</i>
	<i>Megalagrion blackburni</i>
	<i>Megalagrion sp.</i>
	<i>Telmatogeton sp.</i>

**Introduced Species**

<b>Amphibians</b>	<i>Bufo marinus</i>
	<i>Rana catesbiana</i>
<b>Crustaceans</b>	<i>Macrobrachium lar</i>
	<i>Procambarus clarkii</i>
<b>Fish</b>	<i>Clarias fuscus</i>
	<i>Poecilia reticulata</i>
	<i>Poecilia sphenops</i>
	Poeciliid sp.
<b>Snails</b>	<i>Pomacea canaliculata</i>

**Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.**

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
<i>Bufo marinus</i>	Introduced	0.25	4	0.4
<i>Rana catesbiana</i>	Introduced	0.25	0.5	0.3
<i>Eleotris sandwicensis</i>	Endemic	1	3.25	1.9
<i>Kuhlia xenura</i>	Endemic	0.75	0.75	0.8
<i>Sicyopterus stimpsoni</i>	Endemic	0.75	3	1.3
<i>Awaous guamensis</i>	Indigenous	0.75	5.5	1.3

<i>Kuhlia</i> sp.	Indigenous	0.5	1.5	1.0
<i>Poecilia sphenops</i>	Introduced	2.5	2.5	2.5
Poeciliid sp.	Introduced	0.25	0.75	0.5

**Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.**

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Eleotris sandwicensis</i>	Endemic		0.48			
<i>Kuhlia xenura</i>	Endemic		1.79			
<i>Sicyopterus stimpsoni</i>	Endemic		1.07			
<i>Awaous guamensis</i>	Indigenous		4.89			
<i>Kuhlia</i> sp.	Indigenous		3.22			
<i>Bufo marinus</i>	Introduced		14.3			
Poeciliid sp.	Introduced		7.04			
<i>Rana catesbiana</i>	Introduced		2.03			

**Species Distributions: Presence (P) of species in different stream reaches.**

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic			P	P	
<i>Eleotris sandwicensis</i>	Endemic		P			
<i>Kuhlia xenura</i>	Endemic		P			
<i>Sicyopterus stimpsoni</i>	Endemic		P	P	P	
<i>Megalagrion blackburni</i>	Endemic			P		
<i>Megalagrion</i> sp.	Endemic			P	P	
<i>Clithon cariosus</i>	Endemic		P			
<i>Neritina granosa</i>	Endemic		P			
<i>Neritina vespertina</i>	Endemic		P			
<i>Awaous guamensis</i>	Indigenous		P	P	P	
Gobiid sp.	Indigenous			P	P	
<i>Kuhlia sandwicensis</i>	Indigenous		P			
<i>Kuhlia</i> sp.	Indigenous		P			
<i>Anax junius</i>	Indigenous		P			
<i>Telmatogeton</i> sp.	Indigenous			P	P	
<i>Bufo marinus</i>	Introduced		P			
<i>Rana catesbiana</i>	Introduced		P			
<i>Macrobrachium lar</i>	Introduced		P			
<i>Procambarus clarkii</i>	Introduced		P			
<i>Clarias fuscus</i>	Introduced		P			
<i>Poecilia reticulata</i>	Introduced		P			
<i>Poecilia sphenops</i>	Introduced		P			
Poeciliid sp.	Introduced		P			

## HISTORIC RANKINGS

**Historic Rankings:** These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Substantial

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

**Current DAR Decision Rule Status:** The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.

Native Insect Diversity  
> 19 spp.

No

Native Macrofauna  
Diversity > 5 spp.

Yes

Absence of Priority 1  
Introduced

No

Abundance of Any  
Native Species

No

Presence of Candidate  
Endangered Species

No

Endangered Newcomb's  
Snail Habitat


No

### CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

#### WATERSHED RATING: Waiehu, Maui


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




### WATERSHED RATING (Cont): Waiehu, Maui


Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.





Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.



### BIOLOGICAL RATING: Waiehu, Maui



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream Rating  
**7**





Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating  
**6**





All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating  
**5**



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.


Stream Rating  
**5**






### OVERALL RATING: Waiehu, Maui

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.



### RATING STRENGTH: Waiehu, Maui

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



### REFERENCES

- 1961. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
- 1978. Timbol, A.S. and Maciolek. Stream Channel Modification in Hawaii. Part A: Statewide Inventory of Streams; Habitat Factors and Associated Biota.
- 1991. Hau, S. Skippy Hau Databook No. 391 Volume 6.
- 1993. Cowie, R.H. Identity, Distribution and Impacts of Introduced Ampullariidae and Viviparidae in the Hawaiian Islands. J. Med & Appl. Malacol., 5. 61-67.
- 1996. Tagawa, A.W. and G.R. Higashi. Management of a Database on the Occurrence, Abundance and Distribution of Native Freshwater Species. Job Progress Report.

2000. Englund, R.A. Report on Aquatic Insect Monitoring of May 2000 in Pelekunu Valley, Moloka'i, Hawai'i. Hawaii Biological Survey.
2004. Benbow, M.E., Burky, A.J., and C.M. Way. Morphological Characteristics and Species Separation of Hawaiian Postlarval Amphidromous Fishes. *Micronesia*, Vol. 37, No. 1. 127-143.
2006. Polhemus, D.A. Maps of Damselfly Locations.
2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.
2008. Hawai'i Division of Aquatic Resources. Impoundment Surveys in DAR Aquatic Surveys Database.