


Hanawī, Maui



WATERSHED FEATURES

Hanawī watershed occurs on the island of Maui. The Hawaiian meaning of the name is unknown. The area of the watershed is 5.7 square mi (14.7 square km), with maximum elevation of 8074 ft (2461 m). The watershed's DAR cluster code is 2, meaning that the watershed is small, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 7% agricultural, 93% conservation, 0% rural, and 0% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other Private</u>
0.0	6.3	83.0	0.0	0.0	0.0	10.7

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

<u>Permanent Biodiversity Protection</u>	<u>Managed for Multiple Uses</u>	<u>Protected but Unmanaged</u>	<u>Unprotected</u>
6.3	83.0	0.0	10.7

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	0.0	0.00	0.00
Low Intensity Developed	0.0	0.00	0.00
Cultivated	0.0	0.00	0.00
Grassland	7.7	0.44	1.14
Scrub/Shrub	2.5	0.14	0.36
Evergreen Forest	87.5	4.98	12.90
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	2.1	0.12	0.31
Unconsolidated Shoreline	0.1	0.00	0.01
Water	0.0	0.00	0.01
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Hanawī is a perennial stream. Total stream length is 14.3 mi (23 km). The terminal stream order is 2.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	0.3	5.8	28.8	65.0

The following stream(s) occur in the watershed:
Hanawī

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1962	1977	1979	1980	1984	1986	1987
1990	1991	1992	2000	2003	2004	

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	3	0	0	4
DAR General Surveys	0	0	0	1	0
DAR Point Quadrat	0	4	19	0	0
HDFG	0	0	3	5	0
Published Report	0	6	17	12	0
Unpublished Report	0	1	0	1	2

BIOTA INFORMATION

Species List

Native Species

Crustaceans	<i>Atyoida bisulcata</i>
Fish	<i>Awaous guamensis</i>
	<i>Eleotris sandwicensis</i>
	Gobiid sp.
	<i>Kuhlia sandwicensis</i>
	<i>Kuhlia</i> sp.
	<i>Kuhlia xenura</i>
	<i>Lentipes concolor</i>
	<i>Sicyopterus stimpsoni</i>
	<i>Stenogobius hawaiiensis</i>
Snails	<i>Neritina granosa</i>

Native Species

Insects	<i>Anax junius</i>
	<i>Anax</i> sp.
	<i>Megalagrion blackburni</i>
	<i>Megalagrion calliphya</i>
	<i>Megalagrion hawaiiense</i>
	<i>Megalagrion nigrohamatum</i>
	<i>nigrohamatum</i>
	<i>Megalagrion nigrohamatum</i>
	<i>nigrolineatum</i>
	<i>Megalagrion pacificum</i>
	<i>Megalagrion</i> sp.
	<i>Procanace</i> sp.
	<i>Telmatogeton</i> sp.

Introduced Species

Amphibians	<i>Rana catesbiana</i>
	<i>Rana rugosa</i>
	Ranid sp.
Crustaceans	<i>Macrobrachium</i> lar
Fish	<i>Oncorhynchus mykiss</i>
Snails	Lymnaeid sp.
	<i>Lymnea</i> sp.
	<i>Oxychilus</i> sp.

Introduced Species

Insects	<i>Cheumatopsyche analis</i>
	Chironomid larvae
	Culicid sp.
	<i>Pantala flavescens</i>

Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
<i>Eleotris sandwicensis</i>	Endemic	2.5	6	4.5
<i>Kuhlia xenura</i>	Endemic	1	3	1.1
<i>Lentipes concolor</i>	Endemic	0.75	2	1.4
<i>Sicyopterus stimpsoni</i>	Endemic	1.5	4	2.3
<i>Awaous guamensis</i>	Indigenous	1	6	2.4

<i>Kuhlia</i> sp.	Indigenous	0.63	1	0.9
<i>Neritina granosa</i>	Endemic	0.06	1.75	0.4

Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Eleotris sandwicensis</i>	Endemic		1.16	0.27		
<i>Kuhlia xenura</i>	Endemic		1.73	1.42		
<i>Lentipes concolor</i>	Endemic			1.51		
<i>Neritina granosa</i>	Endemic		1.73	21.1		
<i>Sicyopterus stimpsoni</i>	Endemic		1.73	2.94		
<i>Awaous guamensis</i>	Indigenous		1.44	1.78		
<i>Kuhlia</i> sp.	Indigenous		1.44	0.27		

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic		P	P	P	
<i>Eleotris sandwicensis</i>	Endemic		P	P		
<i>Kuhlia xenura</i>	Endemic		P	P		
<i>Lentipes concolor</i>	Endemic		P	P	P	
<i>Sicyopterus stimpsoni</i>	Endemic		P	P		
<i>Megalagrion blackburni</i>	Endemic		P			
<i>Megalagrion calliphya</i>	Endemic		P			P
<i>Megalagrion hawaiiense</i>	Endemic					P
<i>Megalagrion nigrohamatum nigrohamatum</i>	Endemic					P
<i>Megalagrion pacificum</i>	Endemic		P			
<i>Megalagrion</i> sp.	Endemic		P	P	P	
<i>Neritina granosa</i>	Endemic		P	P		
<i>Awaous guamensis</i>	Indigenous		P	P		
Gobiid sp.	Indigenous			P		
<i>Kuhlia sandwicensis</i>	Indigenous		P			
<i>Kuhlia</i> sp.	Indigenous		P	P		
<i>Anax junius</i>	Indigenous		P	P	P	
<i>Anax</i> sp.	Indigenous			P		
<i>Telmatogeton</i> sp.	Indigenous				P	
<i>Telmatogeton</i> sp.	Indigenous				P	
<i>Rana catesbiana</i>	Introduced				P	
<i>Rana rugosa</i>	Introduced				P	
Ranid sp.	Introduced		P		P	
<i>Macrobrachium</i> lar	Introduced		P	P		
<i>Oncorhynchus mykiss</i>	Introduced				P	
<i>Cheumatopsyche analis</i>	Introduced		P	P		
Chironomid larvae	Introduced				P	

Culicid sp.	Introduced	P	P	
<i>Pantala flavescens</i>	Introduced	P	P	P
Lymnaeid sp.	Introduced		P	P
<i>Lymnea</i> sp.	Introduced			P

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Outstanding

U.S. Fish and Wildlife Service High Quality Stream (1988): Yes

The Nature Conservancy- Priority Aquatic Sites (1985): Yes

National Park Service - Nationwide Rivers Inventory (1982): Yes

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity <u>> 19 spp.</u>	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 <u>Introduced</u>
No	Yes	No
Abundance of Any <u>Native Species</u>	Presence of Candidate <u>Endangered Species</u>	Endangered Newcomb's <u>Snail Habitat</u>
No	Yes	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Hanawī, Maui


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.





Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.



WATERSHED RATING (Cont): Hanawā, Maui



Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.

Watershed
Rating
2





Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.

Watershed
Rating
8





Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.

Watershed
Rating
5



Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.

Watershed
Rating
8





BIOLOGICAL RATING: Hanawī, Maui

Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream
Rating



9



Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream
Rating



9



All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream
Rating



8



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.

Stream
Rating

8




OVERALL RATING: Hanawī, Maui

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.

Watershed
Rating


9

**RATING STRENGTH: Hanawī, Maui**

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.

Information
Rating

8

**REFERENCES**

1961. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
1977. Maciolek, J.A. Taxonomic Status, Biology, and Distribution of Hawaiian Lentipes, a Diadromous Goby.
1979. Ford, J.I. Biology of a Hawaiian Fluvial Gastropod *Neritina granosa* Sowerby (Prosobranchia: Neritidae). Masters Thesis.
1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution and Relative Abundance of the Endemic Freshwater Goby, *Lentipes concolor* in Hawaii. Hawaii Cooperative Fishery Research Unit.

1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution, Relative Abundance, and Stream Environment of *Lentipes concolor* (Gill, 1860), and Associated Fauna in Hawaiian Streams.
1984. Kinzie III, R.A, Ford, J.I., Yuen, A.R. and S.J.L. Chow. Habitat Utilization Curves for Native Hawaiian Streams Fishes. Special Report No. 6:20:84.
1986. Kinzie III, R.A.I, Ford, J.I., Yuen, A.R. and S.J.L. Chow. Habitat Modeling of Hawaiian Streams. Technical Report No. 171.
1988. Kinzie III, R.A. Habitat Utilization by Hawaiian Stream Fishes with Reference to Community Structure in Oceanic Island Streams. *Environmental Biology of Fishes*, Vol. 22, No. 3. 179-192.
1990. Hau, S.. Memo Notebook: 1990 FW Stream Surveys Vol. 2.
1991. Brasher, A.M. Baseline Aquatic Survey of Kahiwa Stream, Makapipi Stream, and Hanawi Stream Maui, Hawaii.
1992. Brasher, A.M. Aquatic Monitoring Survey of Hanawi Stream, Makapipi Stream and Palauhulu Stream Maui, Hawaii.
1992. Unknown. Makamaka‘ole Stream Fisheries Survey-Draft Report.
1993. Englund, R.A. and D.A. Polhemus. A survey of the fish and aquatic insect fauna of the Hanawi and Makamakaole Streams Maui, Hawai‘i.
1997. Brasher, A.M. Aquatic Monitoring Survey of Hanawi Stream, Makapipi Stream and Palauhulu Stream Maui, Hawaii.
2000. Dendel. P. Aquatic Monitoring Survey of Hanawi Stream, Makapipi Stream and Palauhulu Stream Maui, Hawaii.
2005. Gingerich, S.B. and R.H. Wolff. Effects of Surface-Water Diversions on Habitat Availability for Native Macrofauna, Northeast Maui, Hawaii.
2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
2008. Hawai‘i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.