


Kawainui, O‘ahu



WATERSHED FEATURES

Kawainui watershed occurs on the island of O‘ahu. The Hawaiian meaning of the name is “the big water”. The area of the watershed is 11.3 square mi (29.2 square km), with maximum elevation of 3081 ft (939 m). The watershed's DAR cluster code is 5, meaning that the watershed is medium size, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 14.2% agricultural, 66.7% conservation, 0% rural, and 19.1% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other Private</u>
0.0	0.1	25.7	0.0	8.7	0.0	65.4

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

<u>Permanent Biodiversity Protection</u>	<u>Managed for Multiple Uses</u>	<u>Protected but Unmanaged</u>	<u>Unprotected</u>
8.7	0.0	25.9	65.4

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	2.6	0.29	0.76
Low Intensity Developed	9.1	1.03	2.67
Cultivated	0.0	0.00	0.00
Grassland	6.5	0.74	1.91
Scrub/Shrub	35.4	3.98	10.32
Evergreen Forest	37.3	4.20	10.89
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	3.9	0.44	1.14
Palustrine Emergent	3.4	0.38	0.99
Estuarine Forested	0.0	0.00	0.00
Bare Land	1.1	0.13	0.33
Unconsolidated Shoreline	0.0	0.00	0.00
Water	0.6	0.07	0.19
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Kawainui is a perennial stream. Total stream length is 22.1 mi (35.6 km). The terminal stream order is 4.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
11.2	0.0	70.5	18.3	0.0

The following stream(s) occur in the watershed:

‘Ainoni	canal	Kahana Iki	Kawainui	Makawao
Maunawili	Olomana	‘Ōma‘o	Palapū	

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1905	1910	1912	1961	1963	1975	1978
1979	1990	1997	1998	1999		

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	1	0	3	4	0
HDFG	2	1	3	0	0
Published Report	1	3	1	0	0
Unpublished Report	2	7	7	0	0

BIOTA INFORMATION

Species List

Native Species

Crustaceans	Amphipod sp. <i>Atyoida bisulcata</i> <i>Macrobrachium grandimanus</i> <i>Macrobrachium sp.</i> <i>Thalamita crenata</i>
Fish	<i>Awaous guamensis</i> <i>Chanos chanos</i> <i>Eleotris sandwicensis</i> <i>Kuhlia sandwicensis</i> <i>Mugil cephalus</i> <i>Sicyopterus stimpsoni</i> <i>Stenogobius hawaiiensis</i>
Worms	Hirudinean sp. <i>Oligochaete sp.</i> <i>Polychaeta sp.</i>

Native Species

Insects	<i>Anax junius</i> <i>Anax sp.</i> <i>Megalagrion hawaiiense</i> <i>Megalagrion nigrohamatum nigrolineatum</i> <i>Megalagrion oceanicum</i> <i>Megalagrion sp.</i> <i>Telmatogeton sp.</i>
----------------	--

Introduced Species

Amphibians	<i>Bufo marinus</i> Ranid sp.
Clams	<i>Corbicula fluminea</i>
Crustaceans	Isopod sp. <i>Macrobrachium lar</i> <i>Procambarus clarkii</i>
Fish	<i>Clarias fuscus</i> <i>Cyprinus carpio</i> <i>Hemichromis elongatus</i> <i>Micropterus dolomieu</i> <i>Micropterus salmoides</i> <i>Poecilia reticulata</i> <i>Tilapia sp.</i> <i>Xiphophorus helleri</i>
Snails	<i>Melania sp.</i> Physid sp. <i>Pomacea canaliculata</i> <i>Pomacea sp.</i>

Introduced Species

Insects	<i>Cheumatopsyche analis</i> Chironomid larvae <i>Ischnura ramburi</i>
----------------	--

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic	P	P	P		
<i>Macrobrachium grandimanus</i>	Endemic		P			
<i>Eleotris sandwicensis</i>	Endemic		P			
<i>Sicyopterus stimpsoni</i>	Endemic	P	P	P		
<i>Stenogobius hawaiiensis</i>	Endemic		P			
<i>Megalagrion hawaiiense</i>	Endemic					P
<i>Megalagrion nigrohamatum</i> <i>nigrolineatum</i>	Endemic			P	P	
<i>Megalagrion oceanicum</i>	Endemic			P	P	
<i>Megalagrion sp.</i>	Endemic	P	P	P		
Amphipod sp.	Indigenous			P		
<i>Thalamita crenata</i>	Indigenous			P		
<i>Awaous guamensis</i>	Indigenous	P	P	P		
<i>Chanos chanos</i>	Indigenous	P	P			
<i>Kuhlia sandwicensis</i>	Indigenous	P	P			
<i>Mugil cephalus</i>	Indigenous	P	P			
<i>Anax junius</i>	Indigenous			P		
<i>Anax sp.</i>	Indigenous	P	P			
<i>Telmatogeton sp.</i>	Indigenous			P		
<i>Bufo marinus</i>	Introduced	P	P	P		
Ranid sp.	Introduced	P	P	P		
Isopod sp.	Introduced	P		P		
<i>Procambarus clarkii</i>	Introduced	P	P	P		
<i>Clarias fuscus</i>	Introduced		P	P		
<i>Cyprinus carpio</i>	Introduced		P			
<i>Micropterus salmoides</i>	Introduced			P		
<i>Poecilia reticulata</i>	Introduced	P	P	P		
<i>Tilapia sp.</i>	Introduced	P	P	P		
<i>Xiphophorus helleri</i>	Introduced	P		P		
<i>Cheumatopsyche analis</i>	Introduced		P	P		
Chironomid larvae	Introduced	P	P	P		
<i>Ischnura ramburi</i>	Introduced	P				
<i>Melania sp.</i>	Introduced	P	P	P		

Physid sp.	Introduced	P	P	
<i>Pomacea canaliculata</i>	Introduced		P	
Hirudinean sp.	Undetermined		P	P
<i>Oligochaete sp.</i>	Undetermined		P	P
<i>Polychaeta sp.</i>	Undetermined			P
<i>Macrobrachium sp.</i>	Unknown	P	P	P

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Moderate

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity <u>> 19 spp.</u>	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 <u>Introduced</u>
No	Yes	No
Abundance of Any <u>Native Species</u>	Presence of Candidate <u>Endangered Species</u>	Endangered Newcomb's <u>Snail Habitat</u>
No	Yes	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Kawainui, O‘ahu


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.





Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.



WATERSHED RATING (Cont): Kawainui, O'ahu



Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.

Watershed
Rating
4





Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.

Watershed
Rating
3





Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.

Watershed
Rating
6





Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.

Watershed
Rating
8





BIOLOGICAL RATING: Kawainui, O'ahu



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream
Rating**7**



Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream
Rating**3**

All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.


Stream
Rating**5**

Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.

Stream
Rating**4**


OVERALL RATING: Kawainui, O‘ahu

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.



RATING STRENGTH: Kawainui, O‘ahu

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

1961. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.

1975. Ford, J.I. Preliminary Biological Survey of Kawainui Marsh and the Maunawili Kahanaki Stream System.

1998. Tagawa, A.W. Management of a Database on the Occurrence, Abundance and Distribution of Native Freshwater Species. Job Progress Report.

1999. Lach, L. and Cowie, R. The Spread of the Introduced Freshwater apple Snail *Pomacea Canaliculata* (Lamarch) (Gastropoda Ampullariidae) on O‘ahu, Hawai‘i. Bishop Museum Occasional Papers: No. 58. 66-71.

2005. Luton, C.D., Brasher, A.M.D., Durkin, C.D. and P. Little. Larval drift of amphidromous shrimps and gobies on the island of O‘ahu, Hawai‘i. *Micronesia* 38 (1). 1-16.
2006. Polhemus, D.A. Maps of Damselfly Locations.
2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
2008. Hawai‘i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.